Crop / Pest	Dosage rate	Remarks
Cotton Red bollworm	15 to 20 kg per ha	Dust plants thoroughly as soon as the pest is observed. Repeat application at 10-to-14-day intervals.
Tobacco African bollworm & Tobacco slug	20 kg per ha	Dust plants thoroughly as soon as the pest is observed. Repeat application when necessary.
Lucerne Lucerne caterpillar	15 to 20 per ha	Dust plants thoroughly as soon as the pest is observed. A single treatment normally gives effective control.
Grape vines Mealybug	20 kg per ha	Only to be used in cases of low infestation pressure. Repeat application at 10-to-14-day intervals.
Lawns Lawn Caterpillars	20 kg / ha OR 20 g / 10 m²	Apply when damage is observed. Irrigate after application.
All Crops and General Garden Pests Elegant Grasshopper	20 kg / ha OR 20 g / 10 m²	Treat areas infested by adult Grasshoppers or Hoppers. A single treatment normally gives effective control.

IT OF REACH OF

Although this remedy has been extensively tested under a large variety of conditions the registration holder does not warrant that it will be efficacious under all conditions because the action and effect thereof may be affected by conditions such as abnormal soil, climatic and storage conditions; compatibility with other substances not indicated on the label and the occurrence of resistance of the pest against the remedy concerned as well as by the method, time and accuracy of application. The registration holder, furthermore, does not accept responsibility for damage to crops, vegetation, the environment, or harm to man or animal or for the lack of performance of the remedy concerned due to failure of the user to

follow the label instructions or to the occurrence of conditions which could not have been foreseen in terms of the registration.

Consult the supplier in the event of any uncertainty.

DIRECTIONS FOR USE: Use only as indicated.

APPLICATION RATES: VETERINARY

AFFLICATION NATES. VETERINARY		
Crop / Pest	Remarks	
Poultry and Cage_ birds Red mites & Tampans	Thoroughly dust all poultry houses, paying attention particularly to the floor, new litter, perches, nests, nest material, as well as cracks in the woodwork and walls.	
Lice	Dust under the wings and vent.	
Dogs and Cats Ticks & Fleas	Dust the hair and rub in well. In the case of cats, brush out the powder afterwards. Treat weekly or as necessary	

Hazard Statements:

Suspected of causing cancer.

Hazardous to the aquatic environment.

Precautionary Statements:

Avoid release into the environment.

Wear impervious rubber gloves and boots, protective clothing and chemical safety googles.



WARNIN



KARBA 50 DP

Reg. No. L1081 - Act No. 36 of 1947

GROUP

KARBA

DUSTING

POWDER

A contact and stomach

for agricultural,

veterinary, home,

and garden uses as listed.

dusting powder insecticide

1A INSECTICIDE

ACTIVE INGREDIENT: carbaryl (carbamate) 50 g/kg

Registered Holder:

(1992/002474/07) PO Box 10413, ASTON MANOR, 1630

Distributed by:

MAKHRO HOME AND GARDEN (PTY) LTD (2014/283523/07) 21 Viben Street, Brackenfell 7560



MEDICAL EMERGENCY NUMBER 0861 555 777

NET VOLUME: 250g